

To appear in ApJ

Dust in an extremely metal-poor galaxy: mid-infrared observations of SBS 0335-052¹

Trinh X. Thuan

Astronomy Department, University of Virginia, P.O. Box 3818, University station,
Charlottesville, VA 22903-0818; txt@virginia.edu

and

Marc Sauvage and Suzanne Madden

CEA/DSM/DAPNIA/Service d'Astrophysique, CE-Saclay, 91191 Gif sur Yvette cedex, France;
msauvage@cea.fr, smadden@cea.fr

ABSTRACT

The metal deficient ($Z = Z_{\odot}/41$) Blue Compact Dwarf Galaxy (BCD) SBS 0335-052 was observed with ISOCAM between 5 and 17 μm . With a $L_{12\mu\text{m}}/L_B$ ratio of 2.15, the galaxy is unexpectedly bright in the mid-infrared for such a low-metallicity object. The mid-infrared spectrum shows no sign of the Unidentified Infrared Bands, which we interpret as an effect of the destruction of their carriers by the very high UV energy density in SBS 0335-052. The spectral energy distribution (SED) is dominated by a very strong continuum which makes the ionic lines of [SiIV] and [NeIII] very weak. From 5 to 17 μm the SED can be fitted with a grey-body spectrum, modified by an extinction law similar to that observed toward the Galactic Center, with an optical depth of $A_V \sim 19\text{--}21$ mag. Such a large optical depth implies that a large fraction (as much as $\sim 75\%$) of the current star-formation activity in SBS 0335-052 is hidden by dust with a mass between $3 \times 10^3 M_{\odot}$ and $5 \times 10^5 M_{\odot}$. Silicate grains are present as silicate extinction bands at 9.7 and 18 μm can account for the unusual shape of the MIR spectrum of SBS 0335-052. It is remarkable that such a nearly primordial environment contains as much dust as galaxies which are 10 times more metal-rich. If the hidden star formation in SBS 0335-052 is typical of young galaxies at high redshifts, then the cosmic star formation rate derived from UV/optical fluxes would be underestimated.

Subject headings: stars: formation – dust, extinction – galaxies: compact – galaxies: individual (SBS 0335-052) – infrared: ISM: continuum – infrared: ISM: lines and bands

¹Based on data obtained with ISO, an ESA project with instruments funded by the ESA member states (especially the PI countries: France, Germany, the Netherlands, and the United Kingdom) with the participation of ISAS and NASA.

1. Introduction

Galaxy formation is one of the most fundamental unsolved problems in astrophysics. Much effort has gone into the search for primeval galaxies (hereafter PGs), i.e. galaxies undergoing their first major burst of star formation, at redshifts larger than ~ 2 . Several objects have been put forward as possible PGs, ranging from high-redshift radio-galaxies to $\text{Ly}\alpha$ emitters (e.g. Steidel et al. 1996; Yee et al. 1996). However, most of these candidate PGs appear to already contain a substantial amount of heavy elements (as indicated for example by the presence of P Cygni profiles), implying previous star formation and metal enrichment.

SBS 0335-052 ($\alpha(1950) = 03^h 35^m 15.2^s$, $\delta(1950) = -05^\circ 12' 25.9''$) is a relatively nearby Blue Compact Dwarf Galaxy (hereafter BCD) with an absolute magnitude $M_B = -16.7$ and which appears to be undergoing its first burst of star formation. With a metallicity of only $Z_\odot/41$ (Izotov et al. 1997), it is the second most metal-deficient galaxy known, after I Zw 18 ($Z_\odot/50$). With a redshift of 0.0136 and a Hubble constant of $75 \text{ km s}^{-1} \text{ Mpc}^{-1}$, it is at a distance of 54.3 Mpc ($1'' = 263 \text{ pc}$). Thuan, Izotov, & Lipovetsky (1997) and Papaderos et al. (1998) have found that the stars in SBS 0335-052 are not older than $\sim 100 \text{ Myr}$, making it a truly young galaxy. Thuan & Izotov (1997) suggest that the large HI envelope within which SBS 0335-052 is embedded may be truly primordial. Yet, despite its youth and extremely low metallicity, HST images of SBS 0335-052 clearly show dust patches mixed in with the six super-star clusters where most of the star formation is occurring (Thuan et al. 1997). The presence of dust, in combination with the intense ultraviolet radiation field from the many young stars in SBS 0335-052 suggests that there may be detectable Mid-Infrared (MIR) emission, as the dust will reprocess the UV starlight and reemit it in the infrared. We have therefore obtained MIR observations of SBS 0335-052 with the Infrared Space Observatory (ISO) to study the properties of dust and star formation in a truly metal-deficient environment, similar to those prevailing at the epoch of galaxy formation.

2. Observations and Data Reduction

The observations were obtained with ISOCAM (Cesarsky et al., 1996a), the MIR imaging instrument aboard ISO (Kessler et al., 1996). They consist of two different sets: a set of broad-band filter maps, and a set obtained by imaging spectroscopy.

The broad-band maps were obtained with a spatial sampling of $3''$ per pixel and in a 3×3 raster mode with a 3-pixel displacement in each direction, giving a total field of view of $114'' \times 114''$. The raster mode increases the sensitivity and improves the flat-field correction since in the center of the map, each sky position is imaged by 9 different camera pixels. The maps were obtained with the LW9, LW10, LW8, LW6 and LW2 filters, centered respectively at 14.9, 12.0, 11.3, 7.7 and $6.7 \mu\text{m}$ (Table 1). Note that the LW10 filter is identical to the IRAS $12 \mu\text{m}$ filter. The integration time was 10.08 s per readout for all filters, except LW10 for which an integration time of 5.04 s was used. Total on-source times were 860 s for LW10 and 1540 s for the other filters.

Imaging spectroscopy was performed using the Circular Variable Filter (CVF) facility of ISOCAM. This mode produces images of the full ISOCAM field of view ($192'' \times 192''$) in the $2\text{--}17\ \mu\text{m}$ wavelength region, with a sampling of $6''$ per pixel and a spectral resolution $\lambda/\Delta\lambda$ of $\simeq 40$. Due to the faintness of the source we obtained observations only in the $8.78\text{--}17.34\ \mu\text{m}$ wavelength range. The individual integration time was 5.04 s and the total on-source time was 4800 s .

The various steps of ISOCAM data reduction are detailed in Stark et al. (1998) and we will only describe here aspects that depart from the methods discussed in that paper: dark current subtraction and transient correction.

The dark current is known to show secular trends which depend on the position of the satellite in its orbit and on the time elapsed since launch. These trends can be accurately modelled (see Biviano et al. 1998), allowing nominal dark subtraction.

Transient corrections using the inversion algorithm of Abergel et al. (1996) were applied successfully to the broad-band observations. For the CVF observations, the known oversimplification of the method combined with the intrinsic faintness of the source results in errors of similar magnitude for both corrected and uncorrected CVF data. The main effect of the correction is to add an artifact at the start of the spectrum (around $17\ \mu\text{m}$) while leaving the rest of the scan unchanged. In particular, synthesis of the LW8 and LW9 flux densities from corrected or uncorrected spectra gives flux densities 1.3 times larger than the observed broad-band flux densities. That the CVF produces higher flux densities than the broad-band filters is to be expected: the operational setup of ISOCAM is such that the scan starts with a short exposure through the LW2 filter with a scale of $6''$ per pixel, giving a much higher illumination on the detector than the scan itself. As the source is very faint, the detector would need a much longer time than our adopted exposure time to stabilize down to its true level. This is not the case for the broad-band maps as the broader filters allow many more photons to reach the detector, thus speeding the stabilization process. As a result, we have adopted for the remaining discussion the CVF spectrum uncorrected for transient effects and scaled down by a factor of 1.3.

Figure 1 displays the spectral energy distribution of SBS 0335-052. The flat-fielding and photometry of the CVF scan were made following Aussel et al. (1998). Conversion from camera units to mJy was performed using the calibration factors given in the *ISOCAM Cookbook*². HST images (Thuan et al. 1997) reveal that most of the star formation in SBS 0335-052 occurs within a region of $\sim 2.5''$ in size, so that the individual super-star clusters in SBS 0335-052 are not resolved by ISOCAM. Thus the flux densities given in Table 1 and plotted in Figure 1 are integrated over the whole star-forming region.

3. Discussion

²available at http://isowww.estec.esa.nl/manuals/iso_cam/.

3.1. Global properties

Since the ISOCAM LW10 filter is equivalent to the IRAS $12\mu\text{m}$ filter, it is interesting to see how the integrated properties of SBS 0335-052 compare with those of other BCDs and irregular galaxies. We have thus compiled IRAS and $\text{H}\alpha$ fluxes for the Thuan & Martin (1981, hereafter TM81) catalog of BCDs, as well as for the sample of BCDs and irregular galaxies in Sauvage, Thuan, & Vigroux (1990) to perform the comparison.

With a $12\mu\text{m}$ to blue luminosity ratio L_{12}/L_B of 2.15 ± 0.06 , SBS 0335-052 is comparable to the most MIR-bright objects in the TM81 catalog. In that sample the mean L_{12}/L_B ratio ($\langle \log L_{12}/L_B \rangle = -0.12 \pm 0.27$), while already twice as high as that for spiral galaxies in the CfA catalog (Sauvage & Thuan 1994), is a factor of 3 smaller than the ratio for SBS 0335-052. This shows that even though the BCD is ~ 10 times more metal-poor than the galaxies in TM81, it nevertheless contains a significant amount of dust heated by an intense UV radiation field.

If we compare the $12\mu\text{m}$ luminosity to a tracer of star formation such as the $\text{H}\alpha$ luminosity (Table 2), SBS 0335-052 appears to be normal. Although total $\text{H}\alpha$ fluxes are scarce for BCDs, the available data give $\langle \log L_{12}/L_{\text{H}\alpha} \rangle = 0.96 \pm 0.70$ for the above samples, as compared to 0.65 ± 0.03 for SBS 0335-052. Thus, even though the metallicity of the BCD is exceptionally low, and the galaxy unusually MIR-bright, its star-forming properties appear to be normal when compared to other BCDs.

3.2. The mid-infrared spectrum

When the MIR spectral energy distribution of SBS 0335-052 (Figure 1) is compared to MIR spectra of other star-forming galaxies (see e.g. Metcalfe et al. 1996; Vigroux 1997), two facts stand out : (1) there is no emission from the so-called Unidentified Infrared Bands (hereafter UIBs, usually attributed to carbon-based dust, see e.g. Papoular et al. 1996; Puget & Léger 1989). In particular the very strong band at $11.3\mu\text{m}$ is completely absent from the spectrum; and (2) there are no evident fine structure ionic lines, even though neon as well as sulfur lines are usually quite bright in starburst galaxies.

In Figure 1, a hint of the $[\text{SiV}]\lambda 10.5\mu\text{m}$ and $[\text{NeIII}]\lambda 15.6\mu\text{m}$ lines can be seen. There is a feature near the position of the $[\text{NeII}]\lambda 12.8\mu\text{m}$ line, but it is centered at $12.8\mu\text{m}$, while the redshifted line falls at $12.98\mu\text{m}$. The quality of our spectrum prevents us from actually measuring line fluxes. Instead we compute upper limits as twice the local rms noise times the instrumental profile width at 20% of peak intensity at the line location. We obtain upper limits of $5.6 \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$ for the $[\text{SiV}]$ line, $4.5 \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$ for the $[\text{NeII}]$ line, and $5.4 \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$ for the $[\text{NeIII}]$ line.

Using the photoionization models of Stasinska (1990) with a metallicity equal to 0.02 that of the sun, and with an integrated $\text{H}\beta$ flux in SBS 0335-052 of $12.1 \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$ (the flux

given by Izotov et al. 1997 was multiplied by 4 to account for aperture and extinction effects as suggested by Thuan & Izotov 1997), we predict the line intensities of [SiIV], [NeIII], and [NeII] to be $5.0 \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$, $1.6 \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$, and $7.7 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ respectively. The predicted intensities are thus completely consistent with the upper limits. We conclude, therefore, that the weakness (or absence) of ionic fine-structure lines in SBS 0335-052 when compared to other star-forming galaxies can be explained by a very strong continuum which decreases the equivalent widths of the lines, making them difficult to detect.

None of the usual UIB features are seen, suggesting that their carbonaceous carriers are absent or have been destroyed. As the star-forming region, 660 pc in diameter, contains 5000 equivalent O7 stars (Izotov et al. 1997), the average energy density is $\gtrsim 10 \text{ eV cm}^{-3}$. Puget & Léger (1989) suggest that at this radiation density, the abundance of the band-emitting dust is reduced by a factor of ~ 10 . Therefore the absence of the bands can be understood as a destruction effect. This is probably enhanced by the very low metallicity of the galaxy which allows harder UV photons to travel further into the interstellar medium.

The origin of the continuum emission is more problematic. We observe what could be interpreted as either a broad emission feature at around $14 \mu\text{m}$, or a broad absorption feature at $\lambda \gtrsim 16 \mu\text{m}$. We favor the second hypothesis as we know of no candidate which would show such an emission feature at $\sim 14 \mu\text{m}$. Very small carbonaceous grains, usually thought to be responsible for the MIR continuum (Désert, Boulanger, & Puget 1990), produce featureless emission. Silicates, the other well-known component of dust, can produce emission features of various shapes, but these are centered around $10 \mu\text{m}$ and even in circumstellar disks never extend beyond $12 \mu\text{m}$ (see e.g. Sitko et al. 1998). On the other hand, silicate extinction bands at 9.7 and $18 \mu\text{m}$ can provide an explanation for the strange shape of the MIR spectrum of SBS 0335-052.

Testing this hypothesis is not straightforward as the expected shape of the dust spectrum in this wavelength range is difficult to constrain. Apart from the stochastic heating of macromolecules which produce the UIBs, MIR emission is thought to be produced by very small grains in a thermal regime intermediate between equilibrium and stochastic heating (see e.g. Désert et al. 1990). We have therefore attempted to fit the spectral energy distribution (SED) of SBS 0335-052 with different screen models where the emission spectrum was successively: (1) a black-body of variable temperature modified by an emissivity law of the form $f_\nu \propto B_\nu(T) \nu^{1-2}$; (2) a power-law spectrum; and (3) the continuum MIR spectrum of the Galactic HII region M17 with UIBs and ionic lines removed (Cesarsky et al. 1996b). The free parameters of the underlying spectra include in all cases a scaling factor, in addition to the temperature in case (1) and the power-law slope in case (2). A further uncertainty comes from the extinction curve to be used. We have tried two different laws, the standard MIR extinction law derived by Draine (1989), and that observed in the direction of the Galactic Center by Lutz et al. (1996). The second law differs from the first by its much higher A_λ/A_V in the range $2\text{--}8 \mu\text{m}$. This difference is probably due to the neglect of ice coating on grains in the Draine (1989) model. The free parameter associated with the extinction curve is simply the absolute extinction at a reference wavelength. Our fitting procedure searches

for best fits to the CVF scan data only. There is no attempt to fit the broad-band data outside of the CVF range. However, a successful model will be one that also accurately predicts them.

Figure 2 shows examples of the three types of fits. Both a power law and the M17 spectrum fail to reproduce the observed SED. Power laws modified by extinction overproduce emission shortward of $8\mu\text{m}$ (Figure 2a) while the M17 spectrum overestimates emission longward of $14\mu\text{m}$ (Figure 2b). On the contrary, a black-body spectrum modified by an emissivity law proportional to $\nu^{1.5}$ and extinguished by a screen of dust, gives an excellent fit to the SED of SBS 0335-052 (Figure 2c). The goodness of the fit depends very little on the exponent of the emissivity law, so its value cannot be used to constrain the nature of the emitting dust. Instead, the fit is sensitive to the shape of the extinction law. It is impossible to obtain a good fit for the LW2 and LW6 bands with the standard extinction curve of Draine (1989). The predicted spectrum always overproduce the emission in these bands by factors of ~ 2 . On the contrary, excellent fits of these bands can be obtained with the Lutz et al. (1996) extinction curve. The temperature obtained for the black-body curve is in the range 240-260 K, but probably carries little physical meaning since the MIR-emitting grains are not likely to be in thermal equilibrium.

The information we derive concerning the dust extinction is more physical. Given the uncertainties in our data, we obtain an equivalent A_V of 19-21 mag. Note that since the screen model maximizes A_V for a given column density of dust, the column density derived from our fit is likely a lower limit to the true column density if emitting and absorbing dust grains are mixed.

Such a high value of extinction is in sharp contrast to the values ($A_V \leq 0.6$ mag) derived by Izotov et al. (1997) from optical spectrophotometric observations. The large difference between the optical and infrared extinctions implies that the heating sources responsible for the infrared emission detected by ISO are probably too deeply embedded in dust to be detected in the visible and that, as in the case of the Antennae galaxies NGC 4038/39 (Mirabel et al. 1998), most of the infrared emission is powered by invisible star clusters. Indeed, correcting the $H\alpha$ flux for 20 mag of visible extinction would lead to unrealistically high star formation rates which would be incompatible with the non-detection of SBS 0335-052 at 1.4 GHz by the NVSS (Condon et al. 1998), or in the far-infrared by IRAS.

The LW10 luminosity of $2.5 \times 10^8 L_\odot$ is equivalent to the bolometric luminosity of 2500 O7 stars, i.e. half the equivalent number of stars in the visible super-star clusters. Given that in the BCD sample of TM81, $\langle \log L_{12}/L_{\text{FIR}} \rangle = -0.85 \pm 0.09$, it is probable that in fact some 18000 equivalent O7 stars are required to power the total infrared emission. This implies that even in very low-metallicity environments such as those characterizing PGs, a significant fraction, i.e. $\sim 3/4$, of the total star formation activity of a galaxy can be effectively hidden from UV/optical observations.

From the extinction derived in the MIR, we can estimate the dust mass. As MIR extinction is essentially due to silicates (Draine & Lee 1984), this result will likely be a lower limit as some carbon-based dust (such as the grains responsible for the continuum) can be present and still

contribute little extinction in the MIR. Using the dust model of Draine & Lee (1984), we obtain a dust surface density of $1.5 \text{ M}_\odot \text{ pc}^{-2}$. To derive a dust mass, we need to estimate lower and upper bounds to the dust spatial extent. For a lower bound, we assume that the dust is only associated with hidden super-star clusters (SSCs). O’Connell, Gallagher, & Hunter (1994) found that SSCs typically have a very compact core with a half-light radius of $\sim 3 \text{ pc}$, embedded in considerably more extended halos with diameters $\sim 30 \text{ pc}$. If we adopt the latter value, and consider that at least 3 SSCs are hidden (they have the power of 2500 equivalent O7 stars needed to power the LW10 luminosity), we derive a dust mass of $\sim 3.2 \times 10^3 \text{ M}_\odot$. This is very probably an underestimate of the true dust mass as there are likely many more hidden SSCs. Furthermore dust is probably not just associated with the SSCs, but rather mixed throughout the star-forming region, especially if it has a supernova origin as argued later on.

A likely upper limit to the spatial extent of the dust is the size of the region where reddening is observed by HST (Thuan et al. 1997), i.e. 660 pc . This gives a dust mass of $5 \times 10^5 \text{ M}_\odot$. With the HI mass given in Table 2, the gas-to-dust mass ratio of SBS 0335-052 is then between $\sim 2 \times 10^3$ and $\sim 3 \times 10^5$, i.e. much higher than in the Galaxy, which is not too surprising for such a low metallicity object (Lisenfeld & Ferrara 1998). More extraordinary is the fact that the dust mass in SBS 0335-052 can be of the same order of magnitude as that in BCDs which are on average ~ 10 times more metal-rich. For comparison $\langle \log M_{\text{dust}} \rangle = 4.4 \pm 0.6 \text{ M}_\odot$ in the TM81 BCD sample.

As discussed by Thuan et al. (1997), because stars in SBS 0335-052 are not older than $\sim 100 \text{ Myr}$, there is not enough time for the silicate dust to be made in the envelopes of red giant stars. Rather silicate grains probably condensed out of the numerous supernova ejecta present in the BCD (see e.g. Lucy et al. (1991) on SN 1987A; Dwek et al. 1992; Wooden et al. 1993). We can check for the plausibility of this hypothesis by adopting, for example, the supernova silicate dust mass input rate of $0.5 \text{ M}_\odot \text{ pc}^{-2} \text{ Gyr}^{-1}$ obtained by Dwek (1998) for the Galactic Center at 0.1 Gyr , the age of SBS 0335-052. Assuming the diameter of the dust-forming region to be 660 pc , we derive a silicate dust mass of $\sim 2 \times 10^4 \text{ M}_\odot$, in the range of the above estimates.

4. Conclusions and Implications

Although SBS 0335-052 is one of the most metal-deficient galaxies known ($Z_\odot/41$), it is unexpectedly bright in the MIR range, implying a silicate dust mass in the range 10^3 – $10^5 \text{ M}_\odot$. The MIR spectrum shows no sign of UIB carriers which are probably destroyed. Despite a difference of a factor of 40 in metallicity, the Galactic Center MIR extinction produces the best fit to the spectrum. This is quite different from the situation in the UV where the extinction law shows a strong dependence on metallicity (Fitzpatrick 1989). This difference is probably due to the fact that the abundance and spectral properties of the carbon-based dust responsible for the UV extinction and UIBs are more metallicity-dependent than those of the silicate grains responsible for the extinction in the MIR. A possible explanation is that the silicate grains are more resistant to photo-destruction than carbon-based dust.

The derived extinction is quite high, $A_V \sim 20$ mag. Given that the total MIR luminosity already requires the bolometric luminosity of 50% more young stars than are seen in the galaxy, the total star formation rate (SFR) as derived from the optical or UV luminosities must underestimate the true star formation rate by at least 50%, and more likely by a factor of 4 as argued above. Thus if SBS 0335-052 is a representative example of PGs, the cosmic star formation rate will be systematically underestimated if based only on UV and optical fluxes. This is in fact the result obtained by Flores et al. (1998) in their ISO survey of distant galaxies. They found that the cosmic SFR derived from FIR luminosities is ~ 2 to 3 times higher than the SFR estimated previously from UV/optical fluxes (Madau et al. 1996).

The ISOCAM data was analyzed using the software package “CIA”, a joint development by the ESA Astrophysics Division and the ISOCAM consortium. The ISOCAM consortium is led by the ISOCAM PI, C. Cesarsky, Direction des Sciences de la Matière, C.E.A, France. T.X.T. has been partially supported by NASA grant JPL961535. We acknowledge useful conversations with Yuri Izotov.

REFERENCES

- Abergel, A., et al. 1996, *A&A*, 315, L329
- Aussel, H., Gerin, M., Boulanger, F., Désert, F. X., Casoli, F., Cutri, R. M., & Signore, M. 1998, *A&A*, 334, L73
- Biviano, A., Sauvage, M., Roman, P., Boulade, O., Gallais, P., & Okumura, K. 1998, “The ISOCAM dark current calibration report”, ESA Technical report
- Cesarsky, C., et al. 1996a, *A&A*, 315, L32
- Cesarsky, D., Lequeux, J., Abergel, A., Perault, M., Palazzi, E., Madden, S., & Tran, D. 1996b, *A&A*, 315, L309
- Condon, J. J., Cotton, W. D., Greisen, E. W., Yin, Q. F., Perley, R. A., Taylor, G. B., & Broderick, J. J. 1998, *AJ*, 115, 1693
- Desert, F. X., Boulanger, F., & Puget, J. L. 1990, *A&A*, 237, 215
- Draine, B. T. 1989, in *Infrared spectroscopy in astronomy*, ed. B. H. Kaldeich, (Noordwijk: ESA), 93
- Draine, B. T., & Lee, H. M. 1984, *ApJ*, 285, 89
- Dwek, E., Moseley, S. H., Glaccum, W., Graham, J. R., Loewenstein, R. F., Silverberg, R. F., & Smith, R. K. 1992, *ApJ*, 389, L21
- Dwek, E. 1998, *ApJ*, 501, 643
- Fitzpatrick, E. L. 1989, in *Interstellar dust*, ed. L. J. Allamandola, & A. G. G. M. Tielens (Dordrecht: Kluwer), 37
- Flores, H., et al. 1998, *ApJ*, submitted
- Izotov, Y., Lipovetsky, V. A., Chaffee, F. H., Foltz, C. B., Guseva, N. G., & Kniazev, A. Y. 1997, *ApJ*, 476, 698
- Kessler, M. F., et al. 1996, *A&A*, 315, L27
- Lisenfeld, U., & Ferrara, A. 1998, *ApJ*, 496, 145
- Lucy, L. B., Danziger, I. J., Gouiffes, C., & Bouchet, P. 1991, in *Supernovae*, ed. S. E. Woosley (New York: Springer), 82
- Lutz, D., et al. 1996, *A&A*, 315, L269
- Madau, P., Ferguson, H., Dickinson, M., Giavalisco, M., Steidel, C., & Fruchter, A. 1996, *MNRAS*, 283, 1388

- Metcalfe, L., et al. 1996, *A&A*, 315, L105
- Mirabel, F., Vigroux, L., Charmandaris, V., Sauvage, M., Gallais, P., Cesarsky, C., Madden, S., & Duc, P.A. 1998, *A&A*, 333, L1
- O’Connell, R. W., Gallagher, J. S., & Hunter, D. A. 1994, *ApJ*, 433, 65
- Papaderos, P., Izotov, Y. I., Fricke, K. J., Thuan, T. X., & Guseva, N. G. 1998, *A&A*, 338, 43
- Papoular, R., Conard, J., Guillois, O., Nenner, I., Reynaud, C., & Rouzaud, J. N. 1996, *A&A*, 315, 222
- Puget, J. L., & Leger, A. 1989, *ARA&A*, 27, 161
- Sauvage, M., & Thuan, T. X. 1994, *ApJ*, 429, 153
- Sauvage, M., Thuan, T. X., & Vigroux, L. 1990, *A&A*, 237, 296
- Sitko, M. L., Grady, C. A., Lynch, D. K., Russel, R. W., & Hanner, M. S. 1998, *ApJ*, submitted
- Starck, J. L., et al. 1998, *A&A*, in press
- Stasinska, G. 1990, *A&AS*, 83, 501
- Steidel, C. C., Giavalisco, M., Pettini, M., Dickinson, M., & Adelberger, K. L. 1996, *ApJ*, 462, L17
- Thuan, T. X., & Martin, G. E. 1981, *ApJ*, 247, 823
- Thuan, T. X., & Izotov, Y. I. 1997, *ApJ*, 489, 623
- Thuan, T. X., Izotov, Y. I., & Lipovetsky, V. A. 1997, *ApJ*, 477, 661
- Thuan, T. X., Lipovetsky, V. A., Martin, J. M., & Pustilnik, S. A. 1998, *A&AS*, in press
- Vigroux, L. 1997, in *Extragalactic astronomy in the infrared*, ed. G. Mamon, T. X. Thuan, & J. T. T. Van, (Paris: Editions Frontières), 63
- Wooden, D. H., Rank, D. M., Bregman, J. D., Witteborn, F. C., Tielens, A. G. G. M., Cohen, M., Pinto, P. A., & Axelrod, T. S. 1993, *ApJS*, 88, 477
- Yee, H. K. C., Ellingson, E., Bechtold, J., Carlberg, R. G., & Cuillandre, J. C. 1996, *AJ*, 111, 1783




Fig. 1.— Spectral energy distribution of SBS 0335-052 in the mid-infrared. The crosses represent broad-band observations at 6.7, 7.7, 11.3, 12.0 and 14.9 μm . The horizontal bars show the width of the broad-band filters. The 1σ errors bars are smaller than the crosses. Broad-band flux densities are derived assuming an intrinsic spectral shape $f_\nu \propto \nu^{-1}$ and integrating over the point spread function. The thick line is the CVF spectrum without transient correction and scaled down by a factor 1.3 (see text for details). The dashed lines show $\pm 1\sigma$ errors. The spectrum has been corrected for effects of the varying point spread function with wavelength. Expected positions of the [SIV] $\lambda 10.5\mu\text{m}$, [NeII] $\lambda 12.8\mu\text{m}$, and [NeIII] $\lambda 15.6\mu\text{m}$ emission lines are indicated.




Fig. 2.— Best-fit models (dashed lines) for the spectral energy distribution of SBS 0335-052. Three different types of emission spectra are considered: (a) a power-law of the form $\nu^{0.7}$ with an extinction $A_V = 27$ mag; (b) the continuum observed in M 17 (Cesarsky et al. 1996b) with $A_V = 0$; (c) a blackbody of temperature 260 K, with an exponent of 1.5 for the emissivity law, and $A_V = 21$ mag. For all three fits the extinction curve of Lutz et al. (1996) gives the best results. To compare our fit to the broad-band data (crosses), we have integrated the fitted spectrum (dashed line) over the band-pass of the ISOCAM filters. The resulting flux densities are shown by open diamonds. Only model (c) is able to reproduce satisfactorily all broad-band observations.

Table 1. MIR flux densities of SBS 0335-052

filter	λ_0^a μm	$\Delta\nu^a$ 10^{12} Hz	$\Delta\lambda^a$ μm	Flux density mJy	1σ mJy	Luminosity ^b $L_\odot$
LW9	14.9	2.29	2.0	35.2	0.9	$7.4 \cdot 10^7$
LW10	12	13.20	7.0	20.5	0.6	$2.5 \cdot 10^8$
LW8	11.3	2.71	1.3	20.7	0.8	$5.2 \cdot 10^7$
LW6	7.7	7.06	1.5	8.9	0.5	$5.8 \cdot 10^7$
LW2	6.7	16.18	3.5	5.5	0.4	$8.2 \cdot 10^7$

^aAssuming a spectral shape $f_\nu \propto \nu^{-1}$, as in the ISOCAM cookbook.

^bAssuming a distance of 54.3 Mpc.

Table 2. Relevant data for SBS 0335-052

Parameter	Value
$L_{H\alpha}^a$	$5.6 \cdot 10^7 L_\odot$
$L_{H\beta}^a$	$2.2 \cdot 10^7 L_\odot$
L_B^b	$1.2 \cdot 10^8 L_\odot$
L_B^c	$7.2 \cdot 10^8 L_B^\odot$
$M_{\text{H I}}^d$	$9.5 \cdot 10^8 M_\odot$

^aFlux in the $1''$ slit of Izotov et al. 1997, corrected for extinction using $A_V = 0.6$ mag, and multiplied by 2 to account for aperture effects as suggested by Thuan & Izotov 1997.

^bComputed from the absolute blue magnitude in Thuan et al. 1997.

^cHere L_B is expressed in units of the solar blue luminosity, where $L_\odot/L_B^\odot \sim 6.25$.

^dThuan et al. 1998.